

What is celiac disease?

- ◆ A malabsorption disease caused by eating gluten - a protein found in foods made with wheat, rye and barley.
- ◆ Causes damage to the small intestine.
- ◆ Can cause food and medications to be absorbed poorly. This can lead to symptoms of starvation, nutrient and medication malabsorption.
- ◆ The only treatment is a strict gluten-free diet. If the diet is followed, the intestinal damage will slowly heal. This can take several months.
- ◆ The disease is lifelong. Intestinal damage occurs each time gluten is consumed.
- ◆ Celiac disease affects one person in every 133 people in the United States.

Diabetes and celiac disease: The link

- ◆ There is a genetic link between Type 1 diabetes and celiac disease.
- ◆ Developing one of the diseases increases the risk of developing the other.
- ◆ The prevalence of celiac disease in people with Type 1 diabetes is about 6% worldwide.
- ◆ When a family has two children with Type 1 diabetes, there is a much higher chance some in the family will have celiac disease.
- ◆ Symptoms of celiac disease vary widely, but are often absent in persons with diabetes.
- ◆ Celiac disease can cause unstable blood sugar control.

Signs of a malabsorption problem

	Diabetes	Celiac Disease
Weight Loss	x	x
GI symptoms (abdominal pain, bloating, cramping, diarrhea, constipation)	x	x
Ketones in urine (from protein wasting)	x	
Less insulin needs (despite large food intake)	x	x
Abnormal labs	x	x

Getting tested:

People with Type 1 diabetes should be tested for celiac disease.

Initial screening should be done within 1 to 2 years of diagnosis of diabetes.

Periodic follow-up screening is advised every few years for persons at risk.

A normal test does not rule out later development of celiac disease - that is why follow-up screening is necessary.

Initial screening is done by a simple blood test.

CELIAC DISEASE

Classic symptoms...

Gas, bloating, diarrhea, constipation, vomiting, weight loss, anemia, chronic fatigue, bone pain, muscle cramps.

Other symptoms...

Balance problems, migraine headaches, seizures, behavior and memory problems, neuropathies, growth and maturation delays, infertility, bone disease, dental enamel defects, and more.

Gluten-free starches

The following grains and starches are allowed on a gluten-free diet:

Amaranth	Nut flours
Arrowroot	Potato
Beans/legumes	Quinoa
Buckwheat	Rice
Corn	Sorghum
Millet	Soy
Montina	Tapioca
	Teff

Foods Containing Gluten

The following foods are not allowed on a gluten-free diet. This is not a complete listing.

Barley	Malt and Malt
Bran	Extract
Bulgar	Rye
Couscous	Semolina
Durum	Spelt
Einkorn	Triticale
Farro	Wheat
Kamut	Wheat Germ
	Wheat Starch

© Copyright Gluten Intolerance Group of North America, 2006. This document may be reproduced for educational purpose only.

Other documents like this one are available at www.GLUTEN.net.

Advances in celiac disease are fast-paced. If this document is more than 2 years old, please visit our Web site for updated documents.

This information should not be used to diagnose or treat celiac disease or diabetes. See your health care team for diagnosis and treatment options specifically for you.

Local Contacts: Red River Celiacs GIG

Stacey Juhnke, Branch Mgr.
R.O.C.K.Coord.
Fargo, ND
701-237-4854

Sandy Kohler
Moorhead, MN
218-233-4898

Susan Frykman
Lakes Area
Elbow Lake, MN
320-528-2404

Deb Zaks
RRC GIG – North
Grand Forks, ND
701- 775-7402

redriverceliacs@gluten.net
www.redriverceliacs.org

Gluten Intolerance Group (GIG) of North America®
31214 124th Ave. S.E.
Auburn, WA 98092-3667
Phone: 253-833-6655
Fax: 253-833-6675
www.GLUTEN.net
info@GLUTEN.net

GIG is a nonprofit 501c3 national organization providing support for per persons with gluten intolerances, in order to live health, productive lives. GIG Branches provide support at a local level.

To make a donation or become a volunteer to GIG, visit our Web site or call the office at 253-833-6655.